

ON CABLED KNOTS, DEHN SURGERY, AND LEFT-ORDERABLE FUNDAMENTAL GROUPS

ADAM CLAY AND LIAM WATSON

ABSTRACT. Previous work of the authors establishes a criterion on the fundamental group of a knot complement that determines when Dehn surgery on the knot will have a fundamental group that is not left-orderable [6]. We provide a refinement of this criterion by introducing the notion of a *decayed* knot; it is shown that Dehn surgery on decayed knots produces surgery manifolds that have non-left-orderable fundamental group for all sufficiently positive surgeries. As an application, we prove that sufficiently positive cables of decayed knots are always decayed knots. These results mirror properties of L-space surgeries in the context of Heegaard Floer homology.

Definition 1. A group G is left-orderable if there exists a partition of the group elements

$$G = \mathcal{P} \sqcup \{1\} \sqcup \mathcal{P}^{-1}$$

satisfying $\mathcal{P} \cdot \mathcal{P} \subseteq \mathcal{P}$ and $\mathcal{P} \neq \emptyset$. The subset $\mathcal{P}$ is called a positive cone.

This is equivalent to G admitting a left-invariant strict total ordering. For background on left-orderable groups relevant to this paper see [2, 6]; a standard reference for the theory of left-orderable groups is [12]. As established by Boyer, Rolfsen and Wiest [2] (compare [11]), the fundamental group $\pi_1(K)$ of the complement of a knot K in S^3 is always left-orderable. Indeed, this follows from the fact that any compact, connected, irreducible, orientable 3-manifold with positive first Betti number has left-orderable fundamental group [2, Theorem 1.1]. However, the question of left-orderability for fundamental groups of rational homology 3-spheres is considerably more subtle (see [2, 6]) and seems closely tied to certain codimension one structures on the 3-manifold (see [2, 3, 17]). Continuing along the lines of [6] this paper focuses on Dehn surgery, an operation on knots that produces rational homology 3-spheres. We recall this construction in order to fix notation and conventions.

For any knot K in S^3 there is a preferred generating set for the peripheral subgroup $\mathbb{Z} \oplus \mathbb{Z} \subset \pi_1(K)$ provided by the knot meridian μ and the Seifert longitude λ . The latter is uniquely determined (up to orientation) by the existence of a Seifert surface for K . We orient μ so that it links positively with K , and orient λ so that $\mu \cdot \lambda = 1$. For any rational number r with reduced form $\frac{p}{q}$ we denote the peripheral element $\mu^p \lambda^q$ by α_r . At the level of the fundamental group, the result of Dehn surgery along α_r is summarized by the short exact sequence

$$1 \rightarrow \langle\langle \alpha_r \rangle\rangle \rightarrow \pi_1(K) \rightarrow \pi_1(S_r^3(K)) \rightarrow 1.$$

Date: March 11, 2011.

Both authors partially supported by NSERC postdoctoral fellowships.

Here $\langle\langle\alpha_r\rangle\rangle$ denotes the normal closure of α_r , and $S_r^3(K)$ is the 3-manifold obtained by attaching a solid torus to the boundary of $S^3 \setminus \nu(K)$, sending the meridian of the torus to a simple closed curve representing the class

$$[\alpha_r] \in H_1(\partial(S^3 \setminus \nu(K)); \mathbb{Z})/\{\pm 1\}.$$

We will blur the distinction between α_r as an element of the fundamental group or as a primitive class in the (projective) first homology of the boundary, and refer to these peripheral elements as slopes.

While many examples of rational homology 3-spheres have left-orderable fundamental group [2], there exist infinite families of knots for which sufficiently positive Dehn surgery (that is, along a slope parametrized by a suitable large rational number) yields a manifold with non-left-orderable fundamental group [6]. To make this precise, consider the set of slopes

$$\mathcal{S}_r = \{\alpha_{r'} \mid r' \geq r\}$$

for some fixed rational r .

Definition 2. *A nontrivial knot K in S^3 is called r -decayed if, for any positive cone $\mathcal{P}$ in $\pi_1(K)$, either $\mathcal{P} \cap \mathcal{S}_r = \mathcal{S}_r$ or $\mathcal{P} \cap \mathcal{S}_r = \emptyset$.*

The existence of decayed knots is established in [6]. For example, the torus knot $T_{p,q}$ is $(pq - 1)$ -decayed (for $p, q > 0$), and the $(-2, 3, q)$ -pretzel knot is $(10 + q)$ -decayed for odd $q \geq 5$ (see Theorem 11). Our interest in this property stems from the following:

Theorem 3. *If K is r -decayed then $\pi_1(S_r^3(K))$ is not left-orderable for all $r' \geq r$.*

As a result, it is not restrictive to assume that r is a positive rational number since $\pi_1(S_0^3(K))$ is always left-orderable [2]. Notice however that it is not immediately clear how Theorem 3 might be applied in practice, as there is no obvious method for checking when a knot is r -decayed. For this reason, in Section 1 we describe an equivalent formulation of r -decay whose statement is more technical, but easier to verify, than the definition. Together with the proof of Theorem 3, the results of Section 1 provide a useful refinement of the ideas from [6].

Results connecting left-orderability and Dehn surgery may be expected to mirror similar results relating to L-spaces, since there is no known example of an L-space with left-orderable fundamental group, while many L-spaces have fundamental group that is not left-orderable. (see [1, 5, 6, 16, 21]). Recall that an L-space is a rational homology sphere with Heegaard Floer homology that is as simple as possible, in the sense that $\text{rk } \widehat{\text{HF}}(Y) = |H_1(Y; \mathbb{Z})|$ (see [15]). Theorem 3 mirrors a fundamental property of knots admitting L-space surgeries: if $S_n^3(K)$ is an L-space, then $S_r^3(K)$ is an L-space as well for any $r \geq n$.

In the interest of further investigating left-orderability of fundamental groups of 3-manifolds along the lines of [6], we consider the behaviour of Dehn surgery on cables of r -decayed knots (for necessary background, see Section 2). Denoting the (p, q) -cable of the knot K as $C_{p,q}(K)$, the main theorem of this article is:

Theorem 4. *If K is r -decayed then $C_{p,q}(K)$ is pq -decayed whenever $\frac{q}{p} > r$.*

The proof of Theorem 4 is contained in Section 2. Notice that combining Theorem 11 and Theorem 4 provides a rather large class of knots for which sufficiently positive surgery yields a non-left-orderable fundamental group.

Dehn surgery on cabled knots and non-left-orderability of the resulting fundamental groups may again be viewed in the context of Heegaard Floer homology. Referring to knots admitting L-space surgeries as L-space knots, Hedden proves:

Theorem 5. [9, Theorem 1.10] *If K is an L-space knot then $C_{p,q}(K)$ is an L-space knot whenever $\frac{q}{p} \geq 2g(K) - 1$.*

Here, the quantity $g(K)$ is the Seifert genus of K . Note that the converse of this statement has been recently established by Hom [10].

In order to assess the strength of Theorem 4, it is natural to ask when Dehn surgery on a cable knot yields a manifold that has left-orderable fundamental group. It turns out that, in the case that K is r -decayed, Theorem 4 is close to describing all possible non-left-orderable surgeries on a cable knot $C_{p,q}(K)$, in the following sense:

Theorem 6. *Suppose that C is the (p, q) -cable of some knot. If $r \in \mathbb{Q}$ satisfies $r < pq - p - q$, then $\pi_1(S_r^3(C))$ is left-orderable.*

This result is a special case of a more general observation pertaining to satellite knots that is discussed in Section 3. Notice that Theorem 6 makes no reference to the original knot being r -decayed. However, restricted to r -decayed knots, Theorem 4 and Theorem 6 combine to produce an interval of surgery coefficients for which the left-orderability of the associated quotient is not determined. More precisely:

Question 7. *If K is r -decayed and C is a (p, q) -cable of K with $\frac{q}{p} > r$, can Theorem 4 and Theorem 6 be sharpened to determine when $\pi_1(S_{r'}^3(C))$ is left-orderable for r' satisfying $pq - p - q < r' \leq pq$?*

Acknowledgments. We thank Josh Greene, Tye Lidman and Dale Rolfsen for helpful comments on an earlier draft of this paper.

1. A PRACTICAL REFORMULATION OF THEOREM 3

We begin with a reformulation of r -decay that will be essential in connecting this work with the results of [6]. This will require the following lemma:

Lemma 8. *Let G be a left-orderable group containing elements g, h . If $g \in \mathcal{P}$ implies $h \in \mathcal{P}$ for every positive cone $\mathcal{P}$, then $g \in \mathcal{P}$ if and only if $h \in \mathcal{P}$.*

Proof. We need only show the converse, namely $h \in \mathcal{P}$ implies $g \in \mathcal{P}$ for every positive cone $\mathcal{P} \subset G$. For a contradiction, suppose this is not the case, so there exists a positive cone such that $h \in \mathcal{P}$ and $g \notin \mathcal{P}$. Consider the positive cone $\mathcal{Q} = \mathcal{P}^{-1}$, defining the reverse ordering of G . This gives $g \in \mathcal{Q}$ and $h \notin \mathcal{Q}$, contradicting our assumption. $\square$

Proposition 9. *A knot K is r -decayed if and only if for every positive cone $\mathcal{P} \subset \pi_1(K)$ there exists a strictly increasing sequence of positive rational numbers $\{r_i\}$ with $r_i \rightarrow \infty$ satisfying*

- (1) $r = r_0$, and
- (2) $\alpha_r \in \mathcal{P}$ implies $\alpha_{r_i} \in \mathcal{P}$ for all i .

Proof. Suppose that K is r -decayed, and let $\mathcal{P}$ be any positive cone. Choose a strictly increasing sequence of rational numbers $\{r_i\}$ with $r_0 = r$ and $r_i \rightarrow \infty$. Whenever $\alpha_r \in \mathcal{P}$ we have $\mathcal{S}_r \cap \mathcal{P} \neq \emptyset$, so that $\mathcal{S}_r \cap \mathcal{P} = \mathcal{S}_r$ since K is r -decayed. It follows that $\alpha_{r_i} \in \mathcal{S}_r \subset \mathcal{P}$ for all i .

To prove the converse, let $\mathcal{P}$ be a positive cone for $\pi_1(K)$. Fix a strictly increasing sequence $\{r_i\}$ of rational numbers limiting to infinity and satisfying (1) and (2). Suppose that $\alpha_r \in \mathcal{P}$, then by assumption $\alpha_{r_i} \in \mathcal{P}$ for all $i > 0$.

Now suppose that $\mu^m \lambda^n$ is an element of $\mathcal{S}_r$. Choose r_i, r_{i+1} with corresponding reduced forms $\frac{p_i}{q_i}, \frac{p_{i+1}}{q_{i+1}}$ such that $r_i < \frac{m}{n} < r_{i+1}$. By solving

$$\begin{aligned} q_i a + q_{i+1} b &= cn \\ p_i a + p_{i+1} b &= cm \end{aligned}$$

we can find positive integers a, b and c such that $(\mu^{p_i} \lambda^{q_i})^a (\mu^{p_{i+1}} \lambda^{q_{i+1}})^b = (\mu^m \lambda^n)^c$. Explicitly, Cramer's rule gives

$$a = \begin{vmatrix} n & q_{i+1} \\ m & p_{i+1} \end{vmatrix}, \quad b = \begin{vmatrix} q_i & n \\ p_i & m \end{vmatrix}, \quad c = \begin{vmatrix} q_i & q_{i+1} \\ p_i & p_{i+1} \end{vmatrix};$$

note that all these quantities are positive because of our restriction $r_i < \frac{m}{n} < r_{i+1}$ (compare [6, Lemma 17]). This shows that $\mu^m \lambda^n$ is positive, since its c -th power is expressed as a product of positive elements. Hence $\mathcal{S}_r \cap \mathcal{P} = \mathcal{S}_r$.

This establishes the implication $\alpha_r \in \mathcal{P} \Rightarrow \mathcal{S}_r \subset \mathcal{P}$ for every positive cone $\mathcal{P}$. By Lemma 8, this is equivalent to $\mathcal{S}_r \cap \mathcal{P} = \mathcal{S}_r$ or $\mathcal{S}_r \cap \mathcal{P} = \emptyset$ for every positive cone $\mathcal{P}$, so that K is r -decayed. $\square$

Remark 10. *In practice, it is often more natural to establish $\alpha_r \in \mathcal{P}$ implies $\alpha_{r_i}^{w_i} \in \mathcal{P}$ for all i , where $w_i \in \mathbb{N}$ (see in particular the proofs of Lemma 13 and Lemma 14). This situation arises when one constructs (for a given positive cone $\mathcal{P}$) a sequence of unreduced rationals $\{r_i\} = \{\frac{p_i}{q_i}\}$ for which $\gcd(p_i, q_i) = w_i \geq 1$, and $\mu^{p_0} \lambda^{q_0} \in \mathcal{P}$ implies $\mu^{p_i} \lambda^{q_i} \in \mathcal{P}$ for all i . Notice that the implication $\alpha_r \in \mathcal{P}$ implies $\alpha_{r_i}^{w_i} \in \mathcal{P}$ still allows us to apply Proposition 9, since $\alpha_{r_i}^{w_i} \in \mathcal{P}$ if and only if $\alpha_{r_i} \in \mathcal{P}$ (this simple observation holds in any left-orderable group). Ultimately, this results in more flexibility in selecting the sequence $\{r_i\}$.*

The equivalence established in Proposition 9 shows that all examples considered in [6] are r -decayed for certain r , as [6, Corollary 11] is a special case of Proposition 9.

Theorem 11. [6, Theorem 24, Theorem 28 and Theorem 30]

- (1) *The (p, q) -torus knot is $(pq - 1)$ -decayed for all positive, relatively prime pairs of integers p, q .*
- (2) *The $(-2, 3, q)$ -pretzel knot is $(10 + q)$ -decayed for all odd $q \geq 5$.*
- (3) *The $(3, q)$ -torus knot with one positive full twist added along two strands is $(3q + 2)$ -decayed, for all positive q congruent to 2 modulo 3.*

Proof. We consider the case of K_q , the $(-2, 3, q)$ -pretzel knot with $q \geq 5$ odd, the other cases are similar. Set $r = 10 + q$, and $r_i = r + i$. It is shown in [6] that for every positive cone $\mathcal{P}$ in $\pi_1(K_q)$, the implication $\alpha_r \in \mathcal{P} \Rightarrow \alpha_{r_i} \in \mathcal{P}$ holds for all $i \geq 0$. This means that for every left-ordering of $\pi_1(K_q)$, the integer sequence $\{r_i\}$ satisfies the properties required by Proposition 9, and we conclude that K_q is r -decayed. $\square$

Note that the above proof illustrates some particularly special behaviour, as the rational sequences $\{r_i\}$ required by Proposition 9 (which *a priori* may be different for each left-ordering) are replaced by a single integer sequence sufficient for every left-ordering. Thus, Proposition 9 provides a more workable method (than used previously) for checking when a knot has surgeries that yield a non-left-orderable fundamental group. Combined with the material established in [6, Section 2], we provide a short proof of Theorem 3.

Proof of Theorem 3. For contradiction, assume that $\pi_1(S_{r'}^3(K))$ is left-orderable for some $r' \geq r$, and consider the short exact sequence

$$1 \rightarrow \langle\langle \alpha_{r'} \rangle\rangle \xrightarrow{i} \pi_1(K) \xrightarrow{f} \pi_1(S_{r'}^3(K)) \rightarrow 1,$$

as defined in the introduction. Let $\mu, \lambda \in \pi_1(K)$ denote the meridian and longitude. Since $\pi_1(S_{r'}^3(K))$ is left-orderable, $\langle\langle \alpha_{r'} \rangle\rangle \cap \langle\mu, \lambda\rangle = \langle\alpha_{r'}\rangle$ (see proof of [6, Proposition 20]). In particular, if we fix an arbitrary rational number $s_0 > r'$, then $f(\alpha_{s_0}) \neq 1$. Thus, we may choose a positive cone $\mathcal{Q}$ in $\pi_1(S_{r'}^3(K))$ that contains $f(\alpha_{s_0})$. Next, choose a positive cone $\mathcal{Q}' \subset \langle\langle \alpha_{r'} \rangle\rangle$ not containing $\alpha_{r'}$, and define a positive cone $\mathcal{P} \subset \pi_1(K)$ by

$$\mathcal{P} = i(\mathcal{Q}') \sqcup f^{-1}(\mathcal{Q}).$$

Note that $\alpha_{r'} \notin \mathcal{P}$, and $\alpha_{s_0} \in \mathcal{P}$.

This is a standard construction for creating a left-ordering of a group using a short exact sequence, here the result is a left-ordering of $\pi_1(K)$ with positive cone $\mathcal{P}$, relative to which the subgroup $\langle\langle \alpha_{r'} \rangle\rangle$ is convex. Because $\langle\langle \alpha_{r'} \rangle\rangle$ is convex, the intersection $\langle\langle \alpha_{r'} \rangle\rangle \cap \langle\mu, \lambda\rangle = \langle\alpha_{r'}\rangle$ is convex in the restriction ordering of $\langle\mu, \lambda\rangle$. Therefore, [6, Proposition 18] shows that all slopes α_s with $s > r'$ must have the same sign. In particular, since α_{s_0} is positive it follows that all slopes α_s with $s > r'$ are positive, so that

$$\mathcal{Q} \cap \mathcal{S}_r = \{\alpha_s | s > r'\}.$$

Therefore, K is not r -decayed. $\square$

We remark that there is a more geometric argument establishing Theorem 3, that relies upon an understanding of the topology of the space of left-orderings of $\mathbb{Z} \oplus \mathbb{Z}$ (see [18, Section 3] and [4, Chapter 6]). Roughly, every left-ordering of the knot group $\pi_1(K)$ restricts to a left-ordering of the peripheral subgroup that defines a line in $\mathbb{Z} \oplus \mathbb{Z}$, with all positive elements of $\mathbb{Z} \oplus \mathbb{Z}$ on one side of the line, and all the negative elements on the other side. As a result, given two rationals $r_1 < r_2$ corresponding to slopes α_{r_1} and α_{r_2} that have the same sign in every left-ordering, no left-ordering can restrict to an ordering of the peripheral subgroup with corresponding slope s between r_1 and r_2 . The proof of Theorem 3 then follows from checking that whenever $\pi_1(S_{r'}^3(K))$ is left-orderable, we can define a left-ordering of $\pi_1(K)$ that restricts to yield a line of slope r' in the peripheral subgroup (compare [6, Proof of Theorem 9]).

2. THE PROOF OF THEOREM 4

We recall the construction of a cabled knot in order to fix notation. Consider the (p, q) -torus knot $T_{p,q}$, where $p, q > 0$ are relatively prime. As the closure of a p -strand braid, this knot may be naturally viewed in a solid torus T by removing a tubular neighbourhood of the braid axis. The complement of $T_{p,q}$ in T is referred to as a (p, q) -cable space. Now given any knot K in S^3 , the cable knot $C_{p,q}(K)$ is obtained by identifying the boundary of T with the boundary of $S^3 \setminus \nu(K)$, identifying the longitude of T with the longitude λ of K . We will denote this cable knot by C whenever this simplified notation does not cause confusion.

The knot group $\pi_1(C)$ may be calculate via the Seifert-Van Kampen Theorem, by viewing the complement $S^3 \setminus \nu(C)$ as the identification of the boundaries of $S^3 \setminus \nu(K)$ and a solid torus $D^2 \times S^1$ along an essential annulus with core curve given by the slope $\mu^q \lambda^p$. If $\pi_1(D^2 \times S^1) = \langle t \rangle$ then this gives rise to a natural amalgamated product

$$\pi_1(C) \cong \pi_1(K) *_{\mu^q \lambda^p = t^p} \mathbb{Z}.$$

Consulting [6, Section 3], the meridian and longitude for C may be calculated as

$$\mu_C = \mu^u \lambda^v t^{-v} \quad \text{and} \quad \lambda_C = \mu_C^{-pq} t^p$$

where u and v are positive integers satisfying $pu - qv = 1$ (compare [20, Proof of Theorem 3.1]).

Suppose that the knot K is r -decayed, and choose cabling coefficients p and q so that $q/p > r$. To begin, we choose a positive cone $\mathcal{P} \subset \pi_1(C)$ and assume that $\mu_C^{pq} \lambda = t^p$ is positive. This means that $t^p = \mu^q \lambda^p \in \mathcal{P}$, so every element $\mu^m \lambda^n$ is positive whenever $m/n > r$, since K is r -decayed.

Our method of proof will be to check that the cable is pq -decayed by using the equivalence from Proposition 9. In particular, we will show that for the given positive cone $\mathcal{P} \subset \pi_1(C)$ there exists an unbounded sequence of increasing rationals $\{r_i\}$ with $r_0 = pq$, such that our assumption $\alpha_{pq} = \mu_C^{pq} \lambda_C \in \mathcal{P}$ implies $\alpha_{r_i} \in \mathcal{P}$ for all $i > 0$.

First consider the case when μ_C is positive in the left-ordering defined by $\mathcal{P}$. Here, $\mu_C^{pq+N} \lambda_C$ is positive for $N \geq 0$, as it is a product of positive elements. Therefore in this case it suffices to choose $r_i = pq + i$ for all $i \geq 0$.

For the remainder of the proof, we assume that μ_C is negative. For repeated use below, we also observe the crucial identity

$$(t^{-v})^p (\mu^u \lambda^v)^p = (t^p)^{-v} \mu^{up} \lambda^{vp} = \mu^{-qv} \lambda^{-pv} \mu^{up} \lambda^{vp} = \mu^{pv-qu} = \mu,$$

and recall that t^p commutes with μ, λ, μ_C , and λ_C . Therefore, we also have

$$(\mu^u \lambda^v)^p (t^{-v})^p = (t^{-v})^p (\mu^u \lambda^v)^p = \mu.$$

Let k be an arbitrary non-negative integer, and consider the element

$$\mu^{-k} (t^{-v} \mu^u \lambda^v) \mu^k.$$

If this element is positive for some k , then the required sequence is provided by Lemma 13 (proved below). Therefore, we may assume that

$$(1) \quad \mu^{-k} (t^{-v} \mu^u \lambda^v) \mu^k \notin \mathcal{P}$$

for all k .

Similarly, for k a non-negative integer, we consider

$$(\mu^{-k} t^{-v} \mu^k)^{p-1} (\mu^u \lambda^v)^{p-1}.$$

If this element is positive for some non-negative k , then we can create the required sequence using Lemma 14 (proved below). Therefore, we may assume that

$$(2) \quad (\mu^{-k} t^{-v} \mu^k)^{p-1} (\mu^u \lambda^v)^{p-1} \notin \mathcal{P}$$

for all k .

Observe that

$$(\mu^{-k} t^{-v} \mu^k)^{p-1} (\mu^u \lambda^v)^{p-1} = (\mu^{-k} t^v \mu^k) (\mu^{-k} t^{-vp} \mu^k) (\mu^{up-u} \lambda^{vp-v}),$$

which, recalling that t^p commutes with the elements μ and λ , simplifies to give

$$(\mu^{-k} t^v \mu^k) (\mu^{-u} \lambda^{-v}) t^{-vp} \mu^{up} \lambda^{vp} = (\mu^{-k} t^v \mu^k) (\mu^{-u} \lambda^{-v}) \mu = \mu^{-k} t^v \lambda^{-v} \mu^{-u} \mu^{k+1} \notin \mathcal{P}$$

for all k . Taking inverses yields

$$\mu^{-k-1} \mu^u \lambda^v t^{-v} \mu^k = \mu^{-k-1} \mu_C \mu^k \in \mathcal{P}.$$

For the following lemmas, let $>$ denote the left-ordering defined by the positive cone $\mathcal{P}$, so that $h > g$ whenever $g^{-1}h \in \mathcal{P}$. We can then calculate:

Lemma 12. *If (1) and (2) hold for all $k \geq 0$, then $\mu^{N+q} \lambda^p$ must be positive for all $N \geq 0$.*

Proof. Since $\mu^{-k-1}\mu_C\mu^k > 1$, left-multiplying by μ^{k+1} gives $\mu_C\mu^k > \mu^{k+1}$ for all $k \geq 0$. Setting $k = 0$ we obtain $\mu_C > \mu$, so that left-multiplying by μ_C gives rise to

$$\mu_C^2 > \mu_C\mu.$$

By setting $k = 1$, we get

$$\mu_C\mu > \mu^2,$$

which combines with the previous expression to give $\mu_C^2 > \mu^2$. Continuing in this manner, we obtain $\mu_C^N > \mu^N$ for all $N \geq 0$. Left-multiplying by t^p , it follows that

$$\mu_C^{N+pq}\lambda_C = \mu_C^N t^p > \mu^N t^p = \mu^{N+q}\lambda^p > 1,$$

where the final inequality follows from the fact that $(N+q)/p > q/p > r$ and K is r -decayed. $\square$

As in the first case, we may now choose the sequence of rationals $r_i = pq + i$ for all $i \geq 0$, and the requirements of Proposition 9 are met.

To conclude the proof, we establish Lemma 13 and Lemma 14.

Lemma 13. *If $\mu^{-k}(t^{-v}\mu^u\lambda^v)\mu^k \in \mathcal{P}$ for some $k \geq 0$, then there exists a sequence of rationals $\{r_i\}$ such that $\alpha_{r_i} > 1$ for all i .*

Proof. For $N \geq 0$, we rewrite μ_C^N as

$$\mu_C^N = \mu^u\lambda^v\mu^k(\mu^{-k}(t^{-v}\mu^u\lambda^v)^N\mu^k)\mu^{-u-k}\lambda^{-v}.$$

Fix a positive integer s that is large enough so that $(sq - u - k)/(sp - v) > r$, this is possible because $q/p > r$. Next, the product $\mu_C^{N+pq}s}\lambda_C^s = \mu_C^N t^{ps}$ becomes $\mu_C^N \mu^{sq}\lambda^{sp}$, which is equal to

$$\mu^{u+k}\lambda^v(\mu^{-k}(t^{-v}\mu^u\lambda^v)^N\mu^k)(\mu^{qs-u-k}\lambda^{ps-v}).$$

This is a product of positive elements, because:

- (1) $\mu^{u+k}\lambda^v > 1$ because $(u+k)/v > q/p > r$, and
- (2) $\mu^{qs-u-k}\lambda^{ps-v} > 1$ because $(sq - u - k)/(sp - v) > r$,

while the quantity $\mu^{-k}(t^{-v}\mu^u\lambda^v)^N\mu^k$ is positive by assumption. Therefore, in this case we choose our sequence of rationals to be

$$r_i = \frac{pq + i}{s}$$

for $i \geq 0$, this guarantees that the associated slopes α_{r_i} are positive in the given left-ordering. $\square$

Lemma 14. *If $\mu^{-k}(t^{-v}\mu^u\lambda^v)\mu^k \notin \mathcal{P}$ for all $k \geq 0$, and $(\mu^{-k}t^{-v}\mu^k)^{p-1}(\mu^u\lambda^v)^{p-1} \in \mathcal{P}$ for some $k \geq 0$, then there exists a sequence of rationals $\{r_i\}$ such that $\alpha_{r_i} > 1$ for all i .*

Proof. Fix $k \geq 0$ such that $\mu^{-k}(t^{-v}\mu^u\lambda^v)\mu^k < 1$ and $(\mu^{-k}t^{-v}\mu^k)^{p-1}(\mu^u\lambda^v)^{p-1} > 1$, and let n be the smallest positive integer such that

$$(\mu^{-k}t^{-v}\mu^k)^n(\mu^u\lambda^v)^n > 1,$$

and

$$(\mu^{-k}t^{-v}\mu^k)^{n-1}(\mu^u\lambda^v)^{n-1} < 1,$$

note that $1 < n \leq q - 1$. Note that we may rearrange these two expressions, so that

$$\mu^{-k}t^{-vn}(\mu^u\lambda^v)^n\mu^k > 1,$$

and

$$\mu^{-k}(\mu^u\lambda^v)^{1-n}t^{-v(1-n)}\mu^k > 1.$$

Then we can rewrite μ_C^N for $N \geq 1$ as follows:

$$\mu_C^N = \mu^{u+k}\lambda^v(\mu^{-k}t^{-v(1-n)}\mu^k)[(\mu^{-k}t^{-vn}(\mu^u\lambda^v)^n\mu^k)\mu^{-k}(\mu^u\lambda^v)^{1-n}t^{-v(1-n)}\mu^k]^{N-1}\mu^{-k}t^{-vn}.$$

In the above expression, the quantity inside the square brackets is a product of positive elements. Denote this quantity by P . Choose an integer s such that $(qs - k)/ps > r$. Then considering the slope $\mu_C^{N+pq(v+s)}\lambda_C^{v+s} = \mu_C^N t^{p(v+s)}$, we find

$$\mu_C^N t^{p(v+s)} = \mu^{u+k}\lambda^v(\mu^{-k}t^{-v(1-n)}\mu^k)P^{N-1}\mu^{qs-k}\lambda^{ps}t^{pv-vn}.$$

This is a product of positive elements, because:

- (1) $\mu^{u+k}\lambda^v > 1$, since $(u+k)/v > q/p > r$.
- (2) $\mu^{-k}t^{-v(1-n)}\mu^k > 1$, because if we consider its p -th power, we can use the fact that t^p commutes with all peripheral elements so that

$$(\mu^{-k}t^{-v(1-n)}\mu^k)^p = t^{-pv(1-n)} > 1.$$

The final inequality follows from $-pv(1-n) > 0$.

- (3) $\mu^{qs-k}\lambda^{ps} > 1$, because s is chosen so that $(qs - k)/ps > r$.
- (4) $t^{pv-vn} > 1$, because $pv - vn > 0$.

Therefore, in this case we may choose our sequence of rationals to be

$$r_i = \frac{i + pq(v+s)}{v+s}$$

for $i \geq 0$, as the corresponding elements $\mu_C^{i+pq(v+s)}\lambda_C^{v+s}$ are positive in the left-ordering for $i \geq 0$. $\square$

3. SURGERY ON SATELLITES

Let T denote the solid torus containing a knot K^P , we require that K^P is not contained in any 3-ball inside T . The knot K^P will be called the pattern knot. Let K^C denote a knot in S^3 , K^C will be called the companion knot. We construct the satellite knot K with pattern K^P and companion K^C as follows.

Let $h : \partial T \rightarrow \partial(S^3 \setminus \nu(K^C))$ denote a diffeomorphism from the boundary of T to the boundary of the complement of $\nu(K^C)$, which carries the longitude of ∂T onto the longitude of the knot K^C . The knot K is then realized as the image of the knot K^P in the manifold

$$S^3 \setminus \nu(K^C) \sqcup_h T = S^3.$$

Lemma 15. [19, Proposition 3.4] *There exists a homomorphism $\phi : \pi_1(K) \rightarrow \pi_1(K^P)$ that preserves peripheral structure.*

Proof. We can compute the fundamental group $\pi_1(K)$ by using the Seifert-Van Kampen theorem. Since

$$S^3 \setminus \nu(K) = S^3 \setminus \nu(K^C) \sqcup_h T \setminus \nu(K^P),$$

the group $\pi_1(K)$ is the free product $\pi_1(K^C) * \pi_1(T \setminus \nu(K^P))$, with amalgamation as follows: The meridian of K^C is identified with the meridian of T , and the longitude of K^C is identified with the longitude of T .

Let N denote the normal closure in $\pi_1(K)$ of the commutator subgroup of $\pi_1(K^C)$. The quotient $\pi_1(K)/N$ can be considered as the result of killing the longitude of T . Topologically we can think of this quotient as gluing a second solid torus T' to the torus T containing K^P , in such a way that the meridian of T' is glued to the longitude of T . The result is that $\pi_1(K^C)$ collapses to a single infinite cyclic subgroup, and the group $\pi_1(K)/N$ is isomorphic to $\pi_1(K^P)$. The desired homomorphism ϕ is the quotient map $\pi_1(K) \rightarrow \pi_1(K)/N$. $\square$

Proposition 16. *Suppose that K is a satellite knot with pattern knot K^P , and $r \in \mathbb{Q}$ is any rational number. If $\pi_1(S_r^3(K^P))$ is left-orderable and $S_r^3(K)$ is irreducible, then $\pi_1(S_r^3(K))$ is left-orderable.*

Proof. By Lemma 15, there exists a homomorphism $\phi : \pi_1(K) \rightarrow \pi_1(K^P)$ that preserves peripheral structure, so there exists an induced map

$$\phi_r : \pi_1(S_r^3(K)) \rightarrow \pi_1(S_r^3(K^P))$$

for every $r \in \mathbb{Q}$. Whenever $\pi_1(S_r^3(K^P))$ is left-orderable the image of ϕ_r is nontrivial and $\pi_1(S_r^3(K))$ is left-orderable [2, Theorem 1.1]. $\square$

Proof of Theorem 6. By [7], pq -surgery on a (p, q) -cable knot yields a reducible manifold. Since the minimal geometric intersection number between reducible slopes is ± 1 [8], r -surgery on a (p, q) -cable yields an irreducible manifold whenever $r < pq - p - q$. Moreover, a (p, q) -cable knot can be described as a satellite knot with pattern knot $T_{p,q}$, the (p, q) -torus knot. Therefore, for $r < pq - p - q$ we can apply Proposition 16 to conclude that $\pi_1(S_r^3(K))$ will be left-orderable whenever $\pi_1(S_r^3(T_{p,q}))$ is left-orderable.

We may now combine known results for surgery on torus knots in this setting. On the one hand, $\pi_1(S_r^3(T_{p,q}))$ is an L-space whenever $r \geq 2g - 1$ [14, Proposition 9.5] (see in particular [9, Lemma 2.13]), where $g = g(T_{p,q})$ is the Seifert genus given by $g(T_{p,q}) = \frac{1}{2}(p-1)(q-1)$. On the other, since $S_r^3(T_{p,q})$ is Seifert fibred or a connect sum of lens spaces for every r [13], $S_r^3(T_{p,q})$ is an L-space if and only if $\pi_1(S_r^3(T_{p,q}))$ is not left-orderable [1] (see also [16, 21]).

In particular, $\pi_1(S_r^3(T_{p,q}))$ is left-orderable whenever r is less than $2g(T_{p,q}) - 1$ and the result follows. $\square$

REFERENCES

- [1] Steven Boyer, Cameron Gordon, and Liam Watson. On L-spaces and left-orderable fundamental groups. Preprint.
- [2] Steven Boyer, Dale Rolfsen, and Bert Wiest. Orderable 3-manifold groups. *Ann. Inst. Fourier (Grenoble)*, 55(1):243–288, 2005.
- [3] Danny Calegari and Nathan M. Dunfield. Laminations and groups of homeomorphisms of the circle. *Invent. Math.*, 152(1):149–204, 2003.
- [4] Adam Clay. *The space of left orderings of a group with applications to topology*. Phd, University of British Columbia, 2010.
- [5] Adam Clay and Dale Rolfsen. Ordered groups, eigenvalues, knots, surgery and L-spaces. Preprint, arXiv:1004.3615.
- [6] Adam Clay and Liam Watson. Left-orderable fundamental groups and Dehn surgery. Preprint, arXiv:1009.4176.
- [7] C. McA. Gordon. Dehn surgery and satellite knots. *Trans. Amer. Math. Soc.*, 275(2):687–708, 1983.
- [8] C. McA. Gordon and J. Luecke. Reducible manifolds and Dehn surgery. *Topology*, 35(2):385–409, 1996.
- [9] Matthew Hedden. On knot Floer homology and cabling. II. *Int. Math. Res. Not. IMRN*, (12):2248–2274, 2009.
- [10] Jennifer Hom. A note on cabling and L-space surgeries. *Algebr. Geom. Topol.*, 11(1):219223, 2011.
- [11] James Howie and Hamish Short. The band-sum problem. *J. London Math. Soc. (2)*, 31(3):571–576, 1985.
- [12] Valeriĭ M. Kopytov and Nikolai Ya. Medvedev. *Right-ordered groups*. Siberian School of Algebra and Logic. Consultants Bureau, New York, 1996.
- [13] Louise Moser. Elementary surgery along a torus knot. *Pacific J. Math.*, 38:737–745, 1971.
- [14] Peter Ozsváth and Zoltán Szabó. Knot Floer homology and rational surgeries. Preprint, arXiv:0504404.
- [15] Peter Ozsváth and Zoltán Szabó. On knot Floer homology and lens space surgeries. *Topology*, 44(6):1281–1300, 2005.
- [16] Thomas Peters. On L-spaces and non left-orderable 3-manifold groups. Preprint, arXiv:0903.4495.
- [17] R. Roberts, J. Shareshian, and M. Stein. Infinitely many hyperbolic 3-manifolds which contain no Reebless foliation. *J. Amer. Math. Soc.*, 16(3):639–679 (electronic), 2003.
- [18] Adam S. Sikora. Topology on the spaces of orderings of groups. *Bull. London Math. Soc.*, 36(4):519–526, 2004.
- [19] Daniel S. Silver and Wilbur Whitten. Knot group epimorphisms. *J. Knot Theory Ramifications*, 15(2):153–166, 2006.
- [20] Chichen M. Tsau. Isomorphisms and peripheral structure of knot groups. *Math. Ann.*, 282(2):343–348, 1988.
- [21] Liam Watson. *Involutions on 3-manifolds and Khovanov homology*. PhD, Université du Québec à Montréal, 2009.

CIRGET, UNIVERSITÉ DU QUÉBEC À MONTRÉAL, CASE POSTALE 8888, SUCCURSALE CENTRE-VILLE, MONTRÉAL QC, H3C 3P8.

E-mail address: aclay@cirget.ca

DEPARTMENT OF MATHEMATICS, UCLA, 520 PORTOLA PLAZA, LOS ANGELES CA, 90095.

E-mail address: lwatson@math.ucla.edu